

Hermaphrodite Orbitals with and without Time-Reversal Symmetry

Raffaele Resta

Dipartimento di Fisica Teorica, Università di Trieste, Italy

The localization tensor, alias second cumulant moment $\langle r_\alpha r_\beta \rangle_c$ of the electron distribution, is an intensive quantity characterizing the ground-state wavefunction, and is finite in any insulator [1]. In absence of time-reversal (TR) symmetry, this same tensor acquires an off-diagonal imaginary part [2], proportional to the Chern number C_1 (in 2d).

I specialize here to the case of noninteracting electrons (either HF or KS). Within either periodic or “open” (cluster-like) boundary conditions we have

$$\text{Re} \langle r_\alpha r_\beta \rangle_c = \frac{1}{2N} \int d\mathbf{r} d\mathbf{r}' (\mathbf{r} - \mathbf{r}')_\alpha (\mathbf{r} - \mathbf{r}')_\beta |\rho(\mathbf{r}, \mathbf{r}')|^2, \quad (1)$$

where ρ is the one-body density matrix, and single orbital occupancy is assumed. When performing a localization transformation upon the occupied orbitals, the real part of $\langle r_\alpha r_\beta \rangle_c$ sets a minimum for the quadratic spread in any given direction, averaged over all the orbitals; this statement holds, again, within both kinds of boundary conditions [3].

From now on, I further specialize to the crystalline case. Then the trace of $\langle r_\alpha r_\beta \rangle_c$ equals the Marzari-Vanderbilt Ω_I (divided by the number of occupied bands) while the orbitals which actually minimize the quadratic spread in a given direction have been called “hermaphrodite orbitals” (Wannier-like in one direction, Bloch-like in the orthogonal ones); they decay faster than any polynomial [4].

As for the cases where TR symmetry is broken, we have some experience only in 2d, for either the quantum-Hall (noninteracting) fluid, or the Haldane model Hamiltonian [5].

In a quantum-Hall fluid the dc longitudinal conductance vanishes, hence the system is effectively an insulator. Then $\text{Re} \langle r_\alpha r_\beta \rangle_c$ is finite, while $\text{Im} \langle r_\alpha r_\beta \rangle_c$ is proportional to C_1 (hence to the Hall conductivity). Despite the finiteness of $\text{Re} \langle r_\alpha r_\beta \rangle_c$, two-dimensional localization of the orbitals *cannot* be achieved. Instead, the hermaphrodite orbitals do exist. All can be worked out analytically in the case of a flat substrate potential, where the density matrix appearing in Eq. (1) has a Gaussian decay and the trace of $\langle r_\alpha r_\beta \rangle_c$ equals precisely the squared magnetic length (at filling one). The hermaphrodite orbitals happen to coincide with the Landau-gauge orbitals [2]; notice that ρ is *not* gauge-invariant by a change of the magnetic gauge (although its modulus is such).

Depending on its parameters, the Haldane model may represent either a normal insulator ($C_1 = 0$) or a Chern insulator ($C_1 = \pm 1$). While the normal case is such by all counts, the Chern case is like the quantum-Hall case: $\text{Re} \langle r_\alpha r_\beta \rangle_c$ is finite, $\text{Im} \langle r_\alpha r_\beta \rangle_c$ is proportional to C_1 , and two-dimensional localization of the orbitals *cannot* be achieved [6]. I conjecture that hermaphrodite orbitals still exist.

Other issues, related to macroscopic orbital magnetization, will be possibly discussed.

[1] R. Resta, J. Phys.: Condens. Matter **14**, R625 (2002).

[2] R. Resta, Phys. Rev. Lett. **95**, 196805 (2005).

[3] R. Resta, J. Chem. Phys. **124**, 104104 (2006).

[4] C. Sgiarovello, M. Peressi, and R. Resta, Phys. Rev. **64**, 115202 (2001).

[5] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).

[6] T. Thonhauser, and D. Vanderbilt, at this workshop.